



NeuroScience Product Guide

	Neurotransmitter									Hormone		Methylation pathway
	Serotonin	GABA	Taurine	Glutamate	PEA	Dopamine	Norepinephrine	Epinephrine	Acetylcholine	Cortisol	Melatonin	
NeuroAdrenal	Calm CP ^{1,4}											
	Key Ingredients: <i>Lagerstroemia speciosa</i> (Banaba leaf) (18% corosolic acid [†]), Phosphatidylserine, Glycine, Taurine									Suggested Use: 2 capsules, 1-2 times daily		
NeuroAdrenal	Calm PRT ^{1,5-7}											
	Key Ingredients: <i>Rhodiola rosea</i> (9-15% rosavins), Phosphatidylserine, Glycine, Taurine									Suggested Use: 3-4 capsules, 1-2 times daily		
Neuro-Enteric	NeuroBiota ^{33,34}											
Key Ingredients: Probiotics									Suggested Use: 1 capsules, 1 time daily			
NeuroImmune	Calm G ⁹⁻¹⁰											
	L-cysteine, L-theanine, Alpha-lipoic acid, CoQ10, L-methionine, 5-MTHF									Suggested Use: 1-2 capsules, 3-4 times daily		
NeuroImmune	Kavinace [®] OS ^{8,13}											
	Key Ingredients: L-theanine, Resveratrol, Astaxanthin, Magnesium, Zinc, Selenium, Melatonin									Suggested Use: 2 capsules at bedtime		
Serotonin/GABA	Alpha GABA ^{15,35-38}											
	Key Ingredients: Passion flower extract (<i>Passiflora incarnate</i>) (arial parts), Lemon balm (<i>Melissa officinalis</i>) root extract, L-taurine, Vitamin B6, L-theanine, Ashwagandha (<i>Withania somnifera</i>) root and leaf extract									Suggested Use: 2-3 capsules, up to 2 times daily		
	Alpha GABA PM ^{11,13-17}											
	Key Ingredients: Valerian (<i>Valeriana officinalis</i>) root extract (Valerenic acids 0.8%), Lemon balm (<i>Melissa officinalis</i>) leaf extract, 5-HTP, L-theanine, Melatonin									Suggested Use: 1-2 capsules, at bedtime		
	GABA Trex ⁸											
	Key Ingredient: L-theanine									Suggested Use: 1-2 chewable tablets, up to 3 times daily		
	Serene ^{9,11,18}											
Key Ingredients: 5-HTP, 5-MTHF, Vitamins B and C									Suggested Use: 1 capsule, 1-2 times daily			
SeroTrex ^{8,11}												
Key Ingredients: L-theanine, 5-HTP									Suggested Use: 1-2 chewable tablets, up to 2 times daily			
TravaCor ^{1,8,9,18,19}												
Key Ingredients: Taurine, L-theanine, 5-HTP, 5-MTHF, Vitamins B and C									Suggested Use: 1-3 capsules, 1-2 times daily			
TravaGen ^{1,8,9,18,19}												
Key Ingredients: L-tryptophan, Taurine, L-theanine, 5-MTHF, Vitamins B and C									Suggested Use: 1-3 capsules, 1-2 times daily			

↑ Mild ↑↑ Moderate ↑↑↑ Strong

The number and direction of the arrows reflect the precursor potential for metabolic end products and/or the mechanisms of action of select ingredient relationships, as compared to other NeuroScience products.

Do not exceed suggested use.

NeuroScience Product Categories ■ Catecholamine ■ NeuroAdrenal ■ NeuroEnteric ■ NeuroImmune ■ Serotonin/GABA

[†] Tryptophan is the precursor to 5-HTP; 5-HTP is the direct precursor to serotonin
[‡] Calm CP 16ct delivers equivalent corosolic acid with Banaba leaf at 2% standardization

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.**



NeuroScience Product Guide

	Neurotransmitter									Hormone	Methylation pathway
	Serotonin	GABA	Taurine	Glutamate	PEA	Dopamine	Norepinephrine	Epinephrine	Acetylcholine	Cortisol	
Catecholamine	AttenTrex^{18,20,39}					↑	↑	↑			
	Key Ingredients: L-tyrosine, Vitamin C, Vitamin B6										
	Suggested Use: 1-2 tablets daily for ages 4-8 years, up to 4 times daily for >18 years										
	Balance D^{9,20,22}						↑↑↑	↑↑	↑		
Key Ingredients: N-acetyl-L-cysteine, N-acetyl-L-tyrosine, <i>Mucuna cochinchinensis</i> (99% L-DOPA), 5-MTHF, Vitamins B and C											
Suggested Use: 1-2 capsules, 1-2 times daily											
Daxitrol Essential^{11,20,25}	↑↑				↑↑	↑↑↑	↑	↑	↑↑		
Key Ingredients: N-acetyl-L-cysteine, L-phenylalanine, Green tea leaf extract (65% EGCG), 5-HTP, <i>Mucuna cochinchinensis</i> (99% L-DOPA), <i>Coleus forskohlii</i> root extract (38% Forskolin), <i>Huperzia serrata</i> aerial parts (1% Huperzine A)											
Suggested Use: 1-3 capsules, up to 3 times daily											
Focus DL^{20,23}					↑↑↑						
Key Ingredient: DL-phenylalanine											
Suggested Use: 1-2 capsules, 1-2 times daily											
NeuroAdrenal	AdreCor^{20,26,27}					↑	↑↑	↑↑		↑	✓
	Key Ingredients: L-methionine, L-tyrosine, L-histidine, <i>Rhodiola rosea</i> (5% rosavins), Green tea leaf extract (65% EGCG), 5-MTHF, Vitamins B and C										
	Suggested Use: 1-3 capsules, 1-2 times daily										
	AdreCor with Licorice Root^{20,26,28,29}					↑	↑	↑		↑↑↑	✓
	Key Ingredients: Glycyrrhizic acid (from Licorice root), L-histidine, L-methionine, L-tyrosine, <i>Rhodiola rosea</i> (5% rosavins), Green tea leaf extract (65% EGCG), 5-MTHF, Vitamins B and C										
Suggested Use: 1-3 capsules, 1-2 times daily											
AdreCor with SAME^{20,26,30}						↑	↑	↑↑↑		↑	✓
Key Ingredients: SAME, L-tyrosine, <i>Rhodiola rosea</i> (5% rosavins), Green tea leaf extract (65% EGCG), 5-MTHF, Vitamins B and C											
Suggested Use: 1 capsule, 1-2 times daily											
ExcitaPlus^{20,22,26,27}						↑↑	↑↑↑	↑		↑	✓
Key Ingredients: L-tyrosine, <i>Rhodiola rosea</i> (5% rosavins), L-methionine, <i>Mucuna cochinchinensis</i> (99% L-DOPA), 5-MTHF, Vitamins B and C											
Suggested Use: 1-2 capsules, 1-2 times daily											
Neuro-Immune									↑↑↑		
Key Ingredients: Acetyl-L-carnitine, Alpha-glycerolphosphorylcholine (50% alpha-GPC), <i>Huperzia serrata</i> aerial parts (1% Huperzine A)											
Suggested Use: 1-3 capsules, 1-2 times daily											

↑ Mild ↑↑ Moderate ↑↑↑ Strong

The number and direction of the arrows reflect the precursor potential for metabolic end products and/or the mechanisms of action of select ingredient relationships, as compared to other NeuroScience products.

Do not exceed suggested use.

For complete product information, please visit www.neuroscienceinc.com

For product references, please see the attached Additional Resources page.



*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

** Methionine is the precursor to SAME; SAME is the direct cofactor needed for the conversion of norepinephrine to epinephrine

(888) 342-7272 www.neuroscienceinc.com

Additional Resources

- Product specification sheets
- NeuroSelect technology
- Digital product catalog
- Support network

NeuroScience embraces the expectations of a rigorously regulated professional grade supplement brand. To learn more about our commitment to quality, visit our website at www.neuroscienceinc.com.

NeuroScience branded ingredients:

AdreCor with SAMe uses Gnosis S.p.A.'s S-Adenosyl-L-Methionine disulfate tosylate (Adomet) protected by U.S. Patent No. US8258115 and US6958233, and (6S)-5-methyltetrahydrofolic acid, glucosamine salt (Quatrefolic®) protected by U.S. Patent No. 7,947,662. Adomet and Quatrefolic are registered trademarks of Gnosis S.p.A. Milan, Italy.

AdreCor, AdreCor with Licorice Root, Balance D, Calm G, ExcitaPlus, Serene, TravaCor, and TravaGen use (6S)-5-methyltetrahydrofolic acid, glucosamine salt (Quatrefolic®) protected by U.S. Patent No. 7,947,662. Quatrefolic is a registered trademark of Gnosis S.p.A. Milan, Italy.

References

1. Kumari, et al. *Adv Exp Med Biol*. 2013;775:19-27.
2. Rollinger J, et al. *Bioorg Med Chem*. 2010;18(4):1507-15.
3. Patel H, et al. *Arabian Journal of Chemistry*. 2015.
4. Judy W, et al. *J Ethnopharmacol*. 2003;87(1):115-7.
5. Hughs JW, et al. *J Psychosomat Res*. 2004;353-58.
6. Nesse RM, et al. *Arch Glen Psyciatr*. 1984;41:771-6.
7. Bystritsky A, et al. *J Alt Comple Med*. 2008;14(2):178-180.
8. Kakuda, et al. *Biosci Biotechnol Biochem*. 2002;66(12):2683-6.
9. Stahl S. *J Clin Psychiatry*. 2008;69(9):1352-3.
10. Berk M, et al. *Trends Pharmacol Sci*. 2013;34(3):167-77.
11. Shaw K, et al. *Cochrane Database Syst Rev*. 2002;(1):CD003198.
12. Usharani P, et al. *Drugs R D*. 2008;9(4):243-50.
13. Pandi-Perumal S, et al. *Prog Neurobiol*. 2008;85(3):225-53.
14. Becker A, et al. *BMC Complement Altern Med*. 2014;14:267.
15. Awad R, et al. *Phytother Res*. 2009;23(8):1075-81.
16. Scaglione F and Zangara A. *J Sleep Discord Manage*. 2017;3:016.
17. Kunz D and Mahlberg R. *J Sleep Res*. 2010;19(4):591-6.
18. May J, et al. *Brain Res Bull*. 2013;90:35-42.
19. Birdsall T. *Altern Med Rev*. 1988;3(4):271-80.
20. Ng J, et al. *Nat Rev Neurol*. 2015; 11:567-84.
21. Tuleun C, et al. *Livestock Res Rural Dev*. 2008;20(10).
22. Pardridge W. *NeuroRx*. 2005;2(1):3-14.
23. Berry. *J Neurochem*. 2004;90(2):257-71.
24. Tun M and Herzon S. *J Exp Pharmacol*. 2012;4:113-23.
25. Xie Z and Miller G. *J Pharmacol Exp Ther*. 2008;352(2):617-28.
26. Duncan T, et al. *Mol Nutr Food Res*. 2013;57(4):628-36.
27. Patak P, et al. *Endocr Res*. 2004;30(4):871-875.
28. van Gelderen C, et al. *Hum Exp Toxicol*. 2000;19(8):434-9.
29. Chapman K, et al. *Physiol Rev*. 2013;93(3):1139-206.
30. Bottiglieri T. *Psychiatr Clin North Am*. 2013;36(1):1-13.
31. Ueland P. *J Inherit Metab Dis*. 2011;34(1):3-15.
32. Abbiati G, et al. *Eur J Drug Metab Pharmacokinet*. 1993;18(2):173-80.
33. Barrett E, et al. *J Appl Microbiol*. 2012;113(2):411-7.
34. Messaoudi M, et al. *Br J Nutr*. 2011;105(5):755-64.
35. Kakuda T, et al. *Biosci Biotechnol Biochem*. 2002;66(12):2683-86.
36. Auddy B, et al. *JANA*. 2008;11(1):50-6.
37. Appel K, et al. *Phytother Res*. 2011;25(6):838-43.
38. Kletke O, et al. *PLoS One*. 2013;8(4):e61733.
39. Dakshinamurti K. *Ann NY Acad Sci*. 1990;585:128-44.

For complete product information, please visit
www.neuroscienceinc.com or call (888) 342-7272

